

NORMAL AUTOMORPHISMS OF A FREE METABELIAN NILPOTENT GROUP

GÉRARD ENDIMIONI

ABSTRACT. An automorphism φ of a group G is said to be normal if $\varphi(H) = H$ for each normal subgroup H of G . These automorphisms form a group containing the group of inner automorphisms. When G is a nonabelian free (or free soluble) group, it is known that these groups of automorphisms coincide, but this is not always true when G is a free metabelian nilpotent group. The aim of this paper is to determine the group of normal automorphisms in this last case.

1. PRELIMINARY RESULTS

In a group G , consider a map $\varphi : G \rightarrow G$ of the form

$$\varphi : x \rightarrow x [x, u_1]^{\lambda(1)} \dots [x, u_m]^{\lambda(m)},$$

where $u_1, \dots, u_m$ are elements of G , the exponents $\lambda(1), \dots, \lambda(m)$ being integers (as usual, the commutator $[a, b]$ is defined by $[a, b] = a^{-1}b^{-1}ab$). When G is metabelian, using the relation $[xy, u] = y^{-1}[x, u]y[y, u]$, it is easy to see that φ is an endomorphism of group. These endomorphisms appear in [7] as a solution to a problem of extension from G' to G for certain endomorphisms (also see [1]).

Now consider in an arbitrary group G the map $\varphi_{u,v} : x \rightarrow x[x, u][x, v]$, with $u, v \in G$ and suppose that $\varphi_{u,v}$ is an endomorphism. It is then easy to show that the equality $\varphi_{u,v}(xy^{-1}) = \varphi_{u,v}(x)\varphi_{u,v}(y^{-1})$ is equivalent to the relation $[u, y][x, v] = [x, v][u, y]$. Consequently, if $\varphi_{u,v}$ is an endomorphism for any $u, v \in G$, the group G is metabelian. One can summarize these preliminary remarks like this.

Proposition 1.1. *In a group G , the following conditions are equivalent:*

2000 *Mathematics Subject Classification.* 20E36, 20F28.

Key words and phrases. Normal Automorphism; Free Metabelian Nilpotent Group.

- (i) every map $\varphi : G \rightarrow G$ of the form $\varphi : x \rightarrow x \prod_{i=1}^m [x, u_i]^{\lambda(i)}$ ($u_i \in G$, $\lambda(i) \in \mathbb{Z}$) is an endomorphism;
(ii) G is metabelian.

It is very easy to see that in an metabelian group, such endomorphisms are not necessarily automorphisms (consider for example the endomorphism $x \rightarrow x[x, b]^{-1}$ in the nonabelian group of order 6 defined by the presentation $\langle a, b \mid a^3 = b^2 = (ab)^2 = 1 \rangle$). But in a nilpotent group, each map of the form

$$x \rightarrow w_0 x^{\lambda(1)} w_1 x^{\lambda(2)} \dots x^{\lambda(n)} w_n \quad (\text{with } \lambda(1) + \lambda(2) + \dots + \lambda(n) = \pm 1)$$

is bijective [2, Theorem 1]. Hence we have:

Proposition 1.2. *In a metabelian nilpotent group G , every map $\varphi : G \rightarrow G$ of the form $\varphi : x \rightarrow x \prod_{i=1}^m [x, u_i]^{\lambda(i)}$ ($u_i \in G$, $\lambda(i) \in \mathbb{Z}$) is an automorphism.*

For convenience sake, in a metabelian nilpotent group, an automorphism of the form $x \rightarrow x \prod_{i=1}^m [x, u_i]^{\lambda(i)}$ will be called a *generalized inner automorphism*. Obviously, every inner automorphism is a generalized inner automorphism, since $u^{-1}xu = x[x, u]$. Notice that in a nilpotent group G of class ≤ 2 , we may write

$$\varphi(x) = x \prod_{i=1}^m [x, u_i]^{\lambda(i)} = x \prod_{i=1}^m [x, u_i^{\lambda(i)}] = x \left[x, \prod_{i=1}^m u_i^{\lambda(i)} \right].$$

Hence $\varphi(x) = x[x, u] = u^{-1}xu$, where $u = \prod_{i=1}^m u_i^{\lambda(i)}$. Consequently, in a nilpotent group of class ≤ 2 , the notions of inner automorphism and generalized inner automorphism coincide. On the other hand, in the free nilpotent group of class 3 and rank 2 freely generated by a and b , it is easy to see that the map $x \rightarrow x[x, a, a]$ is a generalized inner automorphism but is not an inner automorphism.

As usual, in a group, the left-normed commutator $[x_1, \dots, x_n]$ is defined inductively by

$$\begin{aligned} [x_1, \dots, x_n] &= [x_1, \dots, x_{n-1}]^{-1} [x_1, \dots, x_{n-1}]^{x_n} \\ &= [x_1, \dots, x_{n-1}]^{-1} x_n^{-1} [x_1, \dots, x_{n-1}] x_n. \end{aligned}$$

The next technical result will be useful in the following.

Proposition 1.3. *In a group G , consider a map $\varphi : G \rightarrow G$ of the form*

$$\varphi(x) = x \prod_{i=1}^n [x, v_{i,1}, \dots, v_{i,\sigma(i)}]^{\eta(i)} \quad (\eta(i) \in \mathbb{Z}),$$

for some function $\sigma : \{1, \dots, n\} \rightarrow \mathbb{N} \setminus \{0\}$ and elements $v_{i,j} \in G$ ($1 \leq i \leq n$, $1 \leq j \leq \sigma(i)$). Then $\varphi(x)$ can be written in the form

$$\varphi(x) = x [x, u_1]^{\lambda(1)} \dots [x, u_m]^{\lambda(m)} \quad (\lambda(i) \in \mathbb{Z}, u_i \in G).$$

Proof. By induction, using the relation $[x, y, z] = [x, y]^{-1} [x, z]^{-1} [x, yz]$ □

Frequently in this paper we shall make use of well-known commutator identities (see for example [10, 5.1.5]). In particular, we have the following relations, valid in a metabelian group G , for any $x, y, z \in G$, $t \in G'$ and $\lambda \in \mathbb{Z}$:

$$\begin{aligned} [xt, y] &= [x, y][t, y] & [t^\lambda, y] &= [t, y]^\lambda \\ [x, y, z][y, z, x][z, x, y] &= 1 & [t, x, y] &= [t, y, x] \end{aligned}$$

Proposition 1.4. *The set of generalized inner automorphisms of a metabelian nilpotent group G forms a (normal) subgroup of the group of automorphisms of G .*

Proof. If φ and ψ are generalized inner automorphisms respectively defined by

$$\varphi(x) = x \prod_{i=1}^m [x, u_i]^{\lambda(i)}, \quad \psi(x) = x \prod_{i=1}^n [x, v_i]^{\mu(i)},$$

an easy calculus shows that

$$\psi \circ \varphi(x) = x \prod_{i=1}^m [x, u_i]^{\lambda(i)} \prod_{i=1}^n [x, v_i]^{\mu(i)} \prod_{i=1}^m \prod_{j=1}^n [x, u_i, v_j]^{\lambda(i)\mu(j)}.$$

Thus $\psi \circ \varphi$ is a generalized inner automorphism by Proposition 1.3. It remains to prove that φ^{-1} is a generalized inner automorphism. For that, it suffices to construct for each integer $k \geq 1$ a generalized inner automorphism ψ_k such that $\psi_k \circ \varphi$ is of the form

$$\psi_k \circ \varphi : x \rightarrow x \prod_{i=1}^m [x, v_{i,1}, \dots, v_{i,\sigma(i)}]^{\eta(i)}$$

for some function $\sigma : \{1, \dots, m\} \rightarrow \mathbb{N} \setminus \{0\}$ and elements $v_{i,j} \in G$ ($1 \leq i \leq m$, $1 \leq j \leq \sigma(i)$), and where each commutator is of weight $\geq 1+2^{k-1}$ (namely, $\sigma(i) \geq 2^{k-1}$ for $i = 1, \dots, m$). Indeed, since G is nilpotent, this implies that $\psi_k \circ \varphi(x) = x$ for k large enough, thus $\varphi^{-1} = \psi_k$ is a generalized inner automorphism, as required. We argue by induction on k . The result is clear when $k = 1$ by taking for ψ_1 the identity map. Now suppose that for some integer $k \geq 1$, there exists a generalized inner automorphism ψ_k such that $\psi_k \circ \varphi(x) = x \prod_{i=1}^m [x, v_{i,1}, \dots, v_{i,\sigma(i)}]^{\eta(i)}$, with $\sigma(i) \geq 2^{k-1}$ for $i = 1, \dots, m$. Put $\psi_{k+1} = \psi' \circ \psi_k$, where ψ' is defined by $\psi'(x) = x \prod_{i=1}^m [x, v_{i,1}, \dots, v_{i,\sigma(i)}]^{-\eta(i)}$. We have

$$\begin{aligned} \psi_{k+1} \circ \varphi(x) &= x \prod_{i=1}^m [x, v_{i,1}, \dots, v_{i,\sigma(i)}]^{\eta(i)} \\ &\times \prod_{j=1}^m \left[x \prod_{i=1}^m [x, v_{i,1}, \dots, v_{i,\sigma(i)}]^{\eta(i)}, v_{j,1}, \dots, v_{j,\sigma(j)} \right]^{-\eta(j)}. \end{aligned}$$

Since

$$\begin{aligned} &\prod_{j=1}^m \left[x \prod_{i=1}^m [x, v_{i,1}, \dots, v_{i,\sigma(i)}]^{\eta(i)}, v_{j,1}, \dots, v_{j,\sigma(j)} \right]^{-\eta(j)} \\ &= \prod_{i=1}^m [x, v_{j,1}, \dots, v_{j,\sigma(j)}]^{-\eta(j)} \prod_{j=1}^m \prod_{i=1}^m [x, v_{i,1}, \dots, v_{i,\sigma(i)}, v_{j,1}, \dots, v_{j,\sigma(j)}]^{-\eta(i)\eta(j)}, \end{aligned}$$

we obtain

$$\psi_{k+1} \circ \varphi(x) = x \prod_{j=1}^m \prod_{i=1}^m [x, v_{i,1}, \dots, v_{i,\sigma(i)}, v_{j,1}, \dots, v_{j,\sigma(j)}]^{-\eta(i)\eta(j)}$$

and this completes the proof of the proposition. $\square$

2. MAIN RESULT

We recall that a normal automorphism φ of a group G is an automorphism such that $\varphi(H) = H$ for each normal subgroup H of G . These automorphisms form a subgroup of the group of all automorphisms of G . Obviously, this subgroup contains the subgroup of inner automorphisms of G . It happens these subgroups coincide, for instance when G is a nonabelian free group [8], a nonabelian free soluble group [11], or a nonabelian free nilpotent group of class 2 [3]. On the other hand,

the subgroup of inner automorphisms is of infinite index in the group of normal automorphisms when G is a nonabelian free nilpotent group of class $k \geq 3$ [3]. Also note there are exactly two normal automorphisms in a (nontrivial) free abelian group: $x \rightarrow x$ and $x \rightarrow x^{-1}$. The determination of all normal automorphisms in a free nilpotent group of class $k \geq 3$ seems to be an open problem. Note however that for $k = 3$, Theorem 2.1 below provides such a determination, since the free nilpotent metabelian group of class 3 (and of given rank) coincide with the corresponding free nilpotent group of class 3. One can find in [5] information about the structure of the group of normal automorphisms of a nilpotent group.

Certainly, in a metabelian nilpotent group, each generalized inner automorphism is a normal automorphism, but a normal automorphism need not to be a generalized inner automorphism. However, our main result states that the converse holds in a nonabelian free metabelian nilpotent group.

Theorem 2.1. *In a nonabelian free metabelian nilpotent group, the group of normal automorphisms coincides with the group of generalized inner automorphisms.*

Before to tackle the proof of this theorem, we give a few consequences. Let φ be an automorphism of a group G . According to Schweigert [12], one says that φ is a polynomial automorphism of G if there exist integers $\epsilon_1, \dots, \epsilon_m \in \mathbb{Z}$ and elements $u_1, \dots, u_m \in G$ such that

$$\varphi(x) = (u_1^{-1}x^{\epsilon_1}u_1) \dots (u_m^{-1}x^{\epsilon_m}u_m)$$

for all $x \in G$. For instance, in a metabelian nilpotent group, any generalized inner automorphism is a polynomial automorphism. It turns out that in a nonabelian free metabelian nilpotent group, these notions coincide.

Corollary 2.1. *An automorphism φ of a nonabelian free metabelian nilpotent group G is a generalized inner automorphism if and only if it is polynomial.*

Proof. Only the part 'if' needs a proof. Therefore, suppose that $\varphi : G \rightarrow G$ is a polynomial automorphism defined by

$$\varphi(x) = (u_1^{-1}x^{\epsilon_1}u_1) \dots (u_m^{-1}x^{\epsilon_m}u_m)$$

and put $\epsilon = \epsilon_1 + \dots + \epsilon_m$. If $\gamma_2(G) = [G', G]$ is the second term of the lower central series of G , it is not difficult to see that φ induces the automorphism $\overline{\varphi} : x \mapsto x^\epsilon[x, u]$ on the free nilpotent group $G/\gamma_2(G)$, where $u = u_1^{\epsilon_1} \dots u_m^{\epsilon_m} \gamma_2(G)$. An easy calculation shows that the relation $\overline{\varphi}(xy) = \overline{\varphi}(x)\overline{\varphi}(y)$ is equivalent to the relation $(xy)^\epsilon = x^\epsilon y^\epsilon$. This implies that $\epsilon = 1$, and so φ^{-1} is polynomial by [2, Theorem 1]. Now it is clear that φ is a normal automorphism, hence φ is a generalized inner automorphism by Theorem 2.1. $\square$

In [4], it is proved that the group generated by all polynomial automorphisms of a metabelian group is itself metabelian. Therefore, we can state:

Corollary 2.2. *The group of normal automorphisms of a free metabelian nilpotent group is metabelian.*

Note that this result is valid even if the relatively free group is abelian, since the group of normal automorphisms is abelian in this case.

Recall that an automorphism of a group G is said to be an IA-automorphism if it induces the identity automorphism on G/G' . We shall write $\text{IA}(G)$ for the group of IA-automorphisms of G . In a nonabelian free metabelian nilpotent group G of class k , the group of normal automorphisms is a subgroup of $\text{IA}(G)$. Since $\text{IA}(G)$ stabilizes the lower central series of G , it is nilpotent of class $k - 1$ (see for instance [13, p. 9]), and thus so is the group of normal automorphisms. Also this result is a consequence of the fact that in a nilpotent group of class $k \geq 2$, the group generated by all polynomial automorphisms is nilpotent of class $k - 1$ [4].

If $d > 1$ denotes the rank of the free metabelian nilpotent group G of class k , note that $\text{IA}(G)$ is metabelian if $d = 2$ (by a result of C.K. Gupta [6]) or if $k \leq 4$ (in this case, $\text{IA}(G)$ is nilpotent of class ≤ 3). On the other hand, it is worth pointing out that contrary to the group of normal automorphisms, $\text{IA}(G)$ is not metabelian if $d > 2$ and $k > 4$. Indeed, without loss of generality, we can assume that G is the free metabelian nilpotent group of class 5 and of rank 3, freely generated by a, b, c . In this group, we define three IA-automorphisms f, g, h by

$$\begin{aligned} f(a) &= a[a, b], \quad f(b) = b, \quad f(c) = c, \\ g(a) &= a, \quad g(b) = b[b, c], \quad g(c) = c, \end{aligned}$$

$$h(a) = a, h(b) = b, h(c) = c[c, a].$$

It follows

$$\begin{aligned} [f, g](a) &= a[c^{-1}, b, a], [f, g](b) = b, [f, g](c) = c, \\ [f, h](a) &= a, [f, h](b) = b, [f, h](c) = c[a, b^{-1}, c], \end{aligned}$$

whence

$$\begin{aligned} [f, h] \circ [f, g](c) &= c[a, b^{-1}, c], \\ [f, g] \circ [f, h](c) &= c[a, b^{-1}, c][c, b, a, b, c]. \end{aligned}$$

Consequently, $[f, g]$ and $[f, h]$ do not commute, hence $\text{IA}(G)$ is not metabelian.

3. PROOF OF THEOREM 2.1.

In all this section, we consider a fixed set S of cardinality ≥ 2 and we denote by M_k the free metabelian nilpotent group of class k freely generated by S . In other words, $M_k = F/F''\gamma_{k+1}(F)$, where F is the free group freely generated by S and $\gamma_{k+1}(F)$ the $(k+1)$ th term of the lower central series of F . The normal closure in a group G of an element a is written $\langle a^G \rangle$.

Lemma 3.1. *For any distinct elements $a, b \in S$ and any integer t , the subgroup $\langle (a^t b)^{M_k} \rangle \cap \gamma_k(M_k) \trianglelefteq M_k$ is generated by the set of elements of the form*

$$[a^t b, c_1, \dots, c_{k-1}], \quad \text{with } c_1, \dots, c_{k-1} \in S \quad (\text{we suppose } k > 1).$$

Moreover, for any subset $\{a = a_0, a_1, \dots, a_r = b\} \subseteq S$ containing a and b , the subgroup

$$\langle (a^t b)^{M_k} \rangle \cap \gamma_k(M_k) \cap \langle a_0, a_1, \dots, a_r \rangle$$

is generated by the set of elements of the form $[a^t b, c_1, \dots, c_{k-1}]$, with $c_1, \dots, c_{k-1} \in \{a_0, a_1, \dots, a_r\}$.

Proof. First suppose that $t = 0$ and consider an element $w \in \langle b^{M_k} \rangle \cap \gamma_k(M_k)$. Hence w is a product of elements of the form $[c_0, c_1, \dots, c_{k-1}]^{\pm 1}$, with $c_i \in S$. More precisely, we can write $w = w_0 w_1$, where w_0 (resp. w_1) is a product of elements of the form $[c_0, c_1, \dots, c_{k-1}]^{\pm 1}$ with $c_i \in S \setminus \{b\}$, (resp. with $c_i \in S$, the element b occurring once at least in $[c_0, c_1, \dots, c_{k-1}]$). In fact, substituting 1 for the indeterminate b in the relation $w = w_0 w_1$ and using the fact that w lies in $\langle b^{M_k} \rangle$, we obtain $w_0 = 1$. Thus w is a product of elements of the

form $[c_0, c_1, \dots, c_{k-1}]^{\pm 1}$, with $c_i = b$ for some $i \in \{0, \dots, k-1\}$. If $i = 1$, we can write $[c_0, b, \dots, c_{k-1}] = [b, c_0, \dots, c_{k-1}]^{-1}$. If $i > 1$, we have $[c_0, c_1, \dots, b, \dots, c_{k-1}] = [c_0, c_1, b, \dots, c_{k-1}]$; by using the relation $[c_0, c_1, b] = [c_1, b, c_0]^{-1}[b, c_0, c_1]^{-1}$, it follows

$$[c_0, c_1, \dots, b, \dots, c_{k-1}] = [b, c_1, c_0, \dots, c_{k-1}][b, c_0, c_1, \dots, c_{k-1}]^{-1}.$$

Thus we have shown that any element of $\langle b^{M_k} \rangle \cap \gamma_k(M_k)$ is a product of elements of the form $[b, c_1, \dots, c_{k-1}]^{\pm 1}$, with $c_i \in S$. Since $[b, c_1, \dots, c_{k-1}] \in \langle b^{M_k} \rangle \cap \gamma_k(M_k)$, the first part of our lemma is proved when $t = 0$.

Now consider the general case. Actually, since clearly $S' = \{a^t b\} \cup S \setminus \{b\}$ is a free basis of M_k , we can use the result obtained in the particular case. It follows that $\langle (a^t b)^{M_k} \rangle \cap \gamma_k(M_k) \leq M_k$ is generated by the set of elements of the form $[a^t b, c_1, \dots, c_{k-1}]$, with $c_i \in S'$. But in fact we may take $c_i \in S$ and so conclude, since

$$[a^t b, c_1, \dots, a^t b, \dots, c_{k-1}] = [a^t b, c_1, \dots, a, \dots, c_{k-1}]^t [a^t b, c_1, \dots, b, \dots, c_{k-1}].$$

Finally, consider an element $w \in \langle (a^t b)^{M_k} \rangle \cap \gamma_k(M_k) \cap \langle a_0, a_1, \dots, a_r \rangle$. we can express w in the form $w = w'w''$, where w' (resp. w'') is a product of elements of the form $[a^t b, c_1, \dots, c_{k-1}]^{\pm 1}$ with $c_1, \dots, c_{k-1} \in \{a_0, a_1, \dots, a_r\}$ (resp. with $c_1, \dots, c_{k-1} \in S$, an element of $S \setminus \{a_0, a_1, \dots, a_r\}$ occurring once at least in the sequence $c_1, \dots, c_{k-1}$). Substituting 1 for all indeterminates in $S \setminus \{a_0, a_1, \dots, a_r\}$, the equality $w = w'w''$ gives $w = w'$. This completes the proof of the lemma. $\square$

As usual, the expression $[x, {}_n y]$ is defined in a group by $[x, {}_0 y] = x$ and $[x, {}_n y] = [[x, {}_{n-1} y], y]$ for each positive integer n .

For a fixed subset $\{a_0, \dots, a_r\} \subseteq S$ and a function $\Delta : \{0, \dots, r\} \rightarrow \mathbb{N}$, we define in M_k the symbol $[x, y, \Delta]$ ($x, y \in M_k$) by

$$[x, y, \Delta] = [x, y, {}_{\Delta(0)} a_0, {}_{\Delta(1)} a_1, \dots, {}_{\Delta(r)} a_r].$$

Note that for any sequence $b_1, \dots, b_k$ of elements of $\{a_0, \dots, a_r\}$, there is a function $\Delta : \{0, \dots, r\} \rightarrow \mathbb{N}$ such that $[x, y, b_1, \dots, b_k] = [x, y, \Delta]$, with $\Delta(0) + \dots + \Delta(r) = k$ (it suffices to apply the relation $[u, v, w] = [u, w, v]$, valid in any metabelian group whenever u belongs to the derived subgroup). If j, j' are distinct given integers in $\{0, \dots, r\}$ and if $\Delta(j) \neq 0$, we define the function $\Delta_{(j)}^{(j')} : \{0, \dots, r\} \rightarrow \mathbb{N}$ by

$$\Delta_{(j)}^{(j')}(j) = \Delta(j) - 1, \quad \Delta_{(j)}^{(j')}(j') = \Delta(j') + 1 \quad \text{and}$$

$$\Delta_{(j)}^{(j')}(i) = \Delta(i) \text{ for all } i \in \{0, \dots, r\} \setminus \{j, j'\}.$$

When Δ is not the null-function, we shall denote by $m(\Delta)$ the least integer j such that $\Delta(j) \neq 0$.

If S is ordered, we may define in M_k *basic commutators* (see for example [9, Chapter 3]. Recall that a basic commutator of weight k' ($2 \leq k' \leq k$) is a commutator of the form $[b_1, b_2, \dots, b_{k'}]$ ($b_i \in S$), with $b_1 > b_2$ and $b_2 \leq b_3 \leq \dots \leq b_{k'}$. Any set of these commutators freely generates a free abelian subgroup of M'_k .

In the next lemma, we aim to express a product of commutateurs of the form $[a_s, a_i, \Delta]$ as a product where only basic commutators occur.

Lemma 3.2. *Let $\{a_0, \dots, a_r\}$ be a finite subset of S ($r > 0$). Choose an integer $s \in \{0, \dots, r\}$ and consider an element $w \in M_{k+2}$ ($k > 0$) of the form*

$$w = \prod_{i, \Delta} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \quad (\epsilon(i, \Delta) \in \mathbb{Z}),$$

where the product is taken over all integers $i \in \{0, \dots, r\}$ and all functions $\Delta : \{0, \dots, r\} \rightarrow \mathbb{N}$ such that $\Delta(0) + \dots + \Delta(r) = k$. Then:

(i) We have

$$\begin{aligned} w &= \prod_{i < s, i \leq m(\Delta)} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \prod_{s < i, i \leq m(\Delta)} [a_i, a_s, \Delta]^{-\epsilon(i, \Delta)} \\ &\times \prod_{s \leq m(\Delta), m(\Delta) < i} [a_i, a_s, \Delta]^{-\epsilon(i, \Delta)} \prod_{m(\Delta) < s, m(\Delta) < i} [a_i, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(s)}]^{-\epsilon(i, \Delta)} \\ &\times \prod_{m(\Delta) < s, m(\Delta) < i} [a_s, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(i)}]^{\epsilon(i, \Delta)} \end{aligned}$$

(in all these products, i lies in $\{0, \dots, r\} \setminus \{s\}$).

(ii) We have $w = 1$ only if all exponents $\epsilon(i, \Delta)$ with $i \in \{0, \dots, r\} \setminus \{s\}$ occuring in the expression of w are zero.

Proof. (i) First we write w as a product of two factors:

$$w = \prod_{i \leq m(\Delta)} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \prod_{m(\Delta) < i} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)}.$$

The first factor can be expressed in the form

$$\begin{aligned} \prod_{i \leq m(\Delta)} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} &= \prod_{i < s, i \leq m(\Delta)} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \prod_{s < i, i \leq m(\Delta)} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \\ &= \prod_{i < s, i \leq m(\Delta)} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \prod_{s < i, i \leq m(\Delta)} [a_i, a_s, \Delta]^{-\epsilon(i, \Delta)}. \end{aligned}$$

Now we deal with the second facteur. We have

$$\begin{aligned} \prod_{m(\Delta) < i} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} &= \prod_{s \leq m(\Delta), m(\Delta) < i} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \prod_{m(\Delta) < s, m(\Delta) < i} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \\ &= \prod_{s \leq m(\Delta), m(\Delta) < i} [a_i, a_s, \Delta]^{-\epsilon(i, \Delta)} \prod_{m(\Delta) < s, m(\Delta) < i} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)}. \end{aligned}$$

Therefore Lemma 3.2(i) is proved if we show the relation

$$(1) \quad \prod_{m(\Delta) < s, m(\Delta) < i} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} = \prod_{m(\Delta) < s, m(\Delta) < i} [a_i, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(s)}]^{-\epsilon(i, \Delta)} \times \prod_{m(\Delta) < s, m(\Delta) < i} [a_s, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(i)}]^{\epsilon(i, \Delta)}.$$

For that, write more explicitly the commutator $[a_s, a_i, \Delta]$ (in the following equalities, we write m instead of $m(\Delta)$):

$$\begin{aligned} [a_s, a_i, \Delta] &= [a_s, a_i, \Delta(0) a_0, \dots, \Delta(r) a_r] \\ &= [a_s, a_i, \Delta(m) a_m, \dots, \Delta(r) a_r] \\ &= [a_s, a_i, a_{m, \Delta(m)-1} a_m, \dots, \Delta(r) a_r]. \end{aligned}$$

Since $[a_s, a_i, a_m] = [a_i, a_m, a_s]^{-1} [a_m, a_s, a_i]^{-1} = [a_i, a_m, a_s]^{-1} [a_s, a_m, a_i]$, we obtain

$$(2) \quad [a_s, a_i, \Delta] = [a_i, a_m, \Delta_{(m)}^{(s)}]^{-1} [a_s, a_m, \Delta_{(m)}^{(i)}].$$

Relation (1) is now an immediate consequence of (2).

(ii) Choose an order in S such that $a_0 < a_1 < \dots < a_r$. Under this condition, all commutators occuring in the expression of w obtained in the first part of the lemma are basic commutators. Suppose that $w = 1$. Since basic commutators of the form $[a_j, a_i, \dots]$ ($i, j \in \{0, \dots, r\} \setminus \{s\}$) occur only in the fourth factor, we have $\epsilon(i, \Delta) = 0$ whenever $m(\Delta) < s$ and $m(\Delta) < i$. It follows

$$\begin{aligned} w &= \prod_{i < s, i \leq m(\Delta)} [a_s, a_i, \Delta]^{\epsilon(i, \Delta)} \prod_{s < i, i \leq m(\Delta)} [a_i, a_s, \Delta]^{-\epsilon(i, \Delta)} \\ &\quad \times \prod_{s \leq m(\Delta), m(\Delta) < i} [a_i, a_s, \Delta]^{-\epsilon(i, \Delta)} = 1. \end{aligned}$$

But all basic commutators occuring in this equality are distinct. Thus $\epsilon(i, \Delta) = 0$ for all integers $i \neq s$ and all functions Δ , as required. $\square$

Lemma 3.3. *Let φ be a normal automorphism of M_{k+2} ($k > 0$) acting trivially on $M_{k+2}/\gamma_{k+2}(M_{k+2})$. Then, for all distinct elements $a, b \in S$, there exists a generalized inner automorphism ψ of M_{k+2} such that $\varphi(a) = \psi(a)$ and $\varphi(b) = \psi(b)$.*

Proof. Let a, b be two distinct elements of S . Then $a^{-1}\varphi(a)$ and $b^{-1}\varphi(b)$ belong to $\langle a^{M_{k+2}} \rangle \cap \gamma_{k+2}(M_{k+2})$ and $\langle b^{M_{k+2}} \rangle \cap \gamma_{k+2}(M_{k+2})$ respectively. By Lemma 3.1, there is a finite subset $\{a = a_0, a_1, \dots, a_r = b\} \subseteq S$ such that

$$\varphi(a) = \varphi(a_0) = a_0 \prod_{i, \Delta} [a_0, a_i, \Delta]^{\alpha(i, \Delta)} \quad (\alpha(i, \Delta) \in \mathbb{Z}),$$

$$\varphi(b) = \varphi(a_r) = a_r \prod_{i, \Delta} [a_r, a_i, \Delta]^{\beta(i, \Delta)} \quad (\beta(i, \Delta) \in \mathbb{Z}),$$

where the two products are taken over all integers $i \in \{0, \dots, r\}$ and all functions $\Delta : \{0, \dots, r\} \rightarrow \mathbb{N}$ with $\Delta(0) + \dots + \Delta(r) = k$ (as in Lemma 3.2, $[a_s, a_i, \Delta]$ is defined by $[a_s, a_i, \Delta] = [a_s, a_i, \Delta(0) a_0, \dots, \Delta(r) a_r]$). Note that if $|S| = 2$ (and so $r = 1$), Lemma 3.3 is easily verified by taking the generalized inner automorphism ψ defined by

$$\psi(x) = x \prod_{\Delta} [x, a_1, \Delta]^{\alpha(1, \Delta)} \prod_{\Delta} [x, a_0, \Delta]^{\beta(0, \Delta)}.$$

Thus we can assume in the following that $|S| > 2$. For any positive integer t , $(a_0^t a_r)^{-1} \varphi(a_0^t a_r) = (a_0^t a_r)^{-1} \varphi(a_0)^t \varphi(a_r)$ belongs to $\langle a_0, a_1, \dots, a_r \rangle$. Therefore, once again by Lemma 2.1, $\varphi(a_0^t a_r)$ can be expressed in the form

$$\varphi(a_0^t a_r) = a_0^t a_r \prod_{i, \Delta} [a_0^t a_r, a_i, \Delta]^{\eta_t(i, \Delta)} \quad (\eta_t(i, \Delta) \in \mathbb{Z}).$$

Since $[a_0, a_i, \Delta]$ is a commutator of weight $k + 2$ in a nilpotent group of class $k + 2$, we obtain

$$\varphi(a_0^t a_r) = a_0^t a_r \prod_{i, \Delta} [a_0, a_i, \Delta]^{t\eta_t(i, \Delta)} \prod_{i, \Delta} [a_r, a_i, \Delta]^{\eta_t(i, \Delta)}.$$

Thus the relation $\varphi(a_0^t a_r) = \varphi(a_0)^t \varphi(a_r)$ implies that

$$\begin{aligned} (3) \quad & \prod_{i, \Delta} [a_0, a_i, \Delta]^{t\eta_t(i, \Delta)} \prod_{i, \Delta} [a_r, a_i, \Delta]^{\eta_t(i, \Delta)} \\ &= \prod_{i, \Delta} [a_0, a_i, \Delta]^{t\alpha(i, \Delta)} \prod_{i, \Delta} [a_r, a_i, \Delta]^{\beta(i, \Delta)}. \end{aligned}$$

Choose an order in S such that $a_0 < a_1 < \dots < a_r$. Then we can use Lemma 3.2(i) (with $s = 0$ or $s = r$) to express each product in (3) as a product of basic commutators (or their inverses). The first product gives

$$\prod_{i,\Delta} [a_0, a_i, \Delta]^{t\eta_t(i,\Delta)} = \prod_{0 < i, i \leq m(\Delta)} [a_i, a_0, \Delta]^{-t\eta_t(i,\Delta)} \prod_{0 \leq m(\Delta), m(\Delta) < i} [a_i, a_0, \Delta]^{-t\eta_t(i,\Delta)}$$

and so $\prod_{i,\Delta} [a_0, a_i, \Delta]^{t\eta_t(i,\Delta)} = \prod_{i \neq 0, \Delta} [a_i, a_0, \Delta]^{-t\eta_t(i,\Delta)}$, where this product is taken over all integers $i \in \{1, \dots, r\}$ and all functions $\Delta : \{0, \dots, r\} \rightarrow \mathbb{N}$ with $\Delta(0) + \dots + \Delta(r) = k$.

Likewise we have $\prod_{i,\Delta} [a_0, a_i, \Delta]^{t\alpha(i,\Delta)} = \prod_{i \neq 0, \Delta} [a_i, a_0, \Delta]^{-t\alpha(i,\Delta)}$.

In the same way, applying Lemma 3.2(i) with $s = r$, the second product of (3) gives

$$\begin{aligned} & \prod_{i,\Delta} [a_r, a_i, \Delta]^{\eta_t(i,\Delta)} = \prod_{i \neq r, i \leq m(\Delta)} [a_r, a_i, \Delta]^{\eta_t(i,\Delta)} \\ & \times \prod_{i \neq r, m(\Delta) < i} [a_i, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(r)}]^{-\eta_t(i,\Delta)} \prod_{i \neq r, m(\Delta) < i} [a_r, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(i)}]^{\eta_t(i,\Delta)}. \end{aligned}$$

Finally, the last product of (3) gives

$$\begin{aligned} & \prod_{i,\Delta} [a_r, a_i, \Delta]^{\beta(i,\Delta)} = \prod_{i \neq r, i \leq m(\Delta)} [a_r, a_i, \Delta]^{\beta(i,\Delta)} \\ & \times \prod_{i \neq r, m(\Delta) < i} [a_i, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(r)}]^{-\beta(i,\Delta)} \prod_{i \neq r, m(\Delta) < i} [a_r, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(i)}]^{\beta(i,\Delta)}. \end{aligned}$$

Thus relation (3) can be written in the form

$$\begin{aligned} (4) \quad & \prod_{i \neq 0, \Delta} [a_i, a_0, \Delta]^{-t\eta_t(i,\Delta)} \prod_{i \neq r, i \leq m(\Delta)} [a_r, a_i, \Delta]^{\eta_t(i,\Delta)} \\ & \times \prod_{i \neq r, m(\Delta) < i} [a_i, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(r)}]^{-\eta_t(i,\Delta)} \prod_{i \neq r, m(\Delta) < i} [a_r, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(i)}]^{\eta_t(i,\Delta)} \\ & = \prod_{i \neq 0, \Delta} [a_i, a_0, \Delta]^{-t\alpha(i,\Delta)} \prod_{i \neq r, i \leq m(\Delta)} [a_r, a_i, \Delta]^{\beta(i,\Delta)} \\ & \times \prod_{i \neq r, m(\Delta) < i} [a_i, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(r)}]^{-\beta(i,\Delta)} \prod_{i \neq r, m(\Delta) < i} [a_r, a_{m(\Delta)}, \Delta_{(m(\Delta))}^{(i)}]^{\beta(i,\Delta)} \end{aligned}$$

(so each commutator occurring in this relation is a basic commutator). Now consider an integer $i \in \{1, \dots, r-1\}$ and a function $\Delta : \{0, \dots, r\} \rightarrow \mathbb{N}$, with $\Delta(0) + \dots + \Delta(r) = k$ (we can always suppose that $r > 1$

since $|S| > 2$). By identifying the exponents of the basic commutator $[a_i, a_0, \Delta]$ of each side of relation (4), it is easy to see that

$$(5) \quad t\eta_t(i, \Delta) + \eta_t\left(i, \Delta_{(r)}^{(0)}\right) = t\alpha(i, \Delta) + \beta\left(i, \Delta_{(r)}^{(0)}\right)$$

if $\Delta(r) > 0$, and $\eta_t(i, \Delta) = \alpha(i, \Delta)$ if $\Delta(r) = 0$. We prove by induction on $\Delta(r)$ that actually, we have always the equality $\eta_t(i, \Delta) = \alpha(i, \Delta)$. At first observe that if $\Delta(r) > 0$, we have $\Delta_{(r)}^{(0)}(r) = \Delta(r) - 1$ and so $\eta_t\left(i, \Delta_{(r)}^{(0)}\right) = \alpha\left(i, \Delta_{(r)}^{(0)}\right)$ by induction. Hence relation (5) implies that

$$\alpha\left(i, \Delta_{(r)}^{(0)}\right) - \beta\left(i, \Delta_{(r)}^{(0)}\right) = t\{\alpha(i, \Delta) - \eta_t(i, \Delta)\}.$$

Consequently, each positive integer t divides the integer $\alpha\left(i, \Delta_{(r)}^{(0)}\right) - \beta\left(i, \Delta_{(r)}^{(0)}\right)$, which is independant of t . It follows that $\alpha\left(i, \Delta_{(r)}^{(0)}\right) = \beta\left(i, \Delta_{(r)}^{(0)}\right)$ and so $\eta_t(i, \Delta) = \alpha(i, \Delta)$, as required.

Using theses relations and taking $t = 1$, relation (3) implies

$$\prod_{\Delta} [a_0, a_r, \Delta]^{\eta(r, \Delta)} \prod_{i, \Delta} [a_r, a_i, \Delta]^{\eta(i, \Delta)} = \prod_{\Delta} [a_0, a_r, \Delta]^{\alpha(r, \Delta)} \prod_{i, \Delta} [a_r, a_i, \Delta]^{\beta(i, \Delta)}$$

(we write η for η_1) and so

$$\prod_{i, \Delta} [a_r, a_i, \Delta]^{\beta(i, \Delta)} = \prod_{i, \Delta} [a_r, a_i, \Delta]^{\eta(i, \Delta)} \prod_{\Delta} [a_r, a_0, \Delta]^{\alpha(r, \Delta) - \eta(r, \Delta)}.$$

Since $\varphi(a_r) = a_r \prod_{i, \Delta} [a_r, a_i, \Delta]^{\beta(i, \Delta)}$, we obtain

$$(6) \quad \varphi(a_r) = a_r \prod_{i, \Delta} [a_r, a_i, \Delta]^{\eta(i, \Delta)} \prod_{\Delta} [a_r, a_0, \Delta]^{\alpha(r, \Delta) - \eta(r, \Delta)}$$

Now consider the generalized inner automorphism ψ defined by

$$\psi(x) = x \prod_{i=1, \dots, r, \Delta} [x, a_i, \Delta]^{\alpha(i, \Delta)} \prod_{\Delta} [x, a_0, \Delta]^{\alpha(r, \Delta) + \eta(0, \Delta) - \eta(r, \Delta)}.$$

We have

$$\psi(a_0) = a_0 \prod_{i=1, \dots, r, \Delta} [a_0, a_i, \Delta]^{\alpha(i, \Delta)} = \varphi(a_0).$$

In the same way,

$$\begin{aligned}
\psi(a_r) &= a_r \prod_{i=1, \dots, r-1, \Delta} [a_r, a_i, \Delta]^{\alpha(i, \Delta)} \prod_{\Delta} [a_r, a_0, \Delta]^{\alpha(r, \Delta) + \eta(0, \Delta) - \eta(r, \Delta)} \\
&= a_r \prod_{i=1, \dots, r-1, \Delta} [a_r, a_i, \Delta]^{\eta(i, \Delta)} \prod_{\Delta} [a_r, a_0, \Delta]^{\alpha(r, \Delta) + \eta(0, \Delta) - \eta(r, \Delta)} \\
&= a_r \prod_{i, \Delta} [a_r, a_i, \Delta]^{\eta(i, \Delta)} \prod_{\Delta} [a_r, a_0, \Delta]^{\alpha(r, \Delta) - \eta(r, \Delta)}
\end{aligned}$$

and so $\psi(a_r) = \varphi(a_r)$ by (6). This completes the proof of Lemma 3.3. $\square$

Lemma 3.3 is actually a weak form of the next result.

Lemma 3.4. *Let φ be a normal automorphism of M_{k+2} ($k > 0$) acting trivially on $M_{k+2}/\gamma_{k+2}(M_{k+2})$. Then φ is a generalized inner automorphism of M_{k+2} .*

Proof. We can assume that $|S| > 2$ (otherwise Lemma 3.4 is a consequence of Lemma 3.3). Consider two distinct elements $a, b \in S$. According to Lemma 3.3, there exists a generalized inner automorphism ψ such that $\varphi(a) = \psi(a)$ and $\varphi(b) = \psi(b)$. It suffices to prove that for any $c \in S \setminus \{a, b\}$, we have $\varphi(c) = \psi(c)$. For that, apply again Lemma 3.3: there are generalized inner automorphisms ψ', ψ'' such that $\varphi(a) = \psi'(a)$, $\varphi(c) = \psi'(c)$ and $\varphi(b) = \psi''(b)$, $\varphi(c) = \psi''(c)$. There exists a finite subset $\{a_0, \dots, a_r\} \subseteq S$, containing a, b, c , such that ψ, ψ', ψ'' can be defined by the equations

$$\begin{aligned}
\psi(x) &= x \prod_{i, \Delta} [x, a_i, \Delta]^{\epsilon(i, \Delta)} \\
\psi'(x) &= x \prod_{i, \Delta} [x, a_i, \Delta]^{\epsilon'(i, \Delta)} \\
\psi''(x) &= x \prod_{i, \Delta} [x, a_i, \Delta]^{\epsilon''(i, \Delta)}
\end{aligned}$$

(the products are taken over all integers $i \in \{0, \dots, r\}$ and all functions $\Delta : \{0, \dots, r\} \rightarrow \mathbb{N}$ with $\Delta(0) + \dots + \Delta(r) = k$). Since $\psi(a) = \psi'(a)$, we have

$$a \prod_{i, \Delta} [a, a_i, \Delta]^{\epsilon(i, \Delta)} = a \prod_{i, \Delta} [a, a_i, \Delta]^{\epsilon'(i, \Delta)}$$

and so

$$\prod_{i, \Delta} [a, a_i, \Delta]^{\epsilon(i, \Delta) - \epsilon'(i, \Delta)} = 1.$$

Applying Lemma 3.2(ii), we obtain $\epsilon(i, \Delta) = \epsilon'(i, \Delta)$ for all functions Δ and all integers $i \in \{0, \dots, r\}$ such that $a_i \neq a$. Similarly, we have $\epsilon(i, \Delta) = \epsilon''(i, \Delta)$ if $a_i \neq b$ and $\epsilon'(i, \Delta) = \epsilon''(i, \Delta)$ if $a_i \neq c$. It follows that $\epsilon(i, \Delta) = \epsilon'(i, \Delta)$ for all function Δ and all integer $i \in \{0, \dots, r\}$, hence $\psi = \psi'$. Thus $\varphi(c) = \psi'(c) = \psi(c)$, as required. $\square$

Proof of Theorem 2.1. We argue by induction on the nilpotency class k of M_k . If $k = 2$, the result follows from [3, Theorem 2(ii)] (in this case, each normal automorphism is inner). Now consider a normal automorphism φ of M_k , with $k > 2$. Then φ induces a normal automorphism on the quotient group $M_k/\gamma_k(M_k)$. By induction, since this quotient is isomorphic to M_{k-1} , there exists a generalized inner automorphism $\psi : M_k \rightarrow M_k$ such that $\varphi(x) = \psi(x)\theta(x)$, where $\theta(x)$ is an element of $\gamma_k(M_k)$. It follows that $\psi^{-1}(\varphi(x)) = x\psi^{-1}(\theta(x))$. Thus $\psi' := \psi^{-1} \circ \varphi$ is a normal automorphism of M_k acting trivially on $M_k/\gamma_k(M_k)$. By Lemma 3.4, ψ' is a generalized inner automorphism, and so is $\varphi = \psi \circ \psi'$. This completes the proof of Theorem 2.1. $\square$

We end with a question: assume φ is a normal automorphism of a free nilpotent group G of class $k \geq 4$ (and of rank > 1). Is φ of the form

$$\varphi : x \rightarrow x [x, u_1]^{\lambda(1)} \dots [x, u_m]^{\lambda(m)} \quad (u_i \in G, \lambda(i) \in \mathbb{Z})?$$

REFERENCES

- [1] A. Caranti and C.M. Scoppola. Endomorphisms of two-generated metabelian groups that induce the identity modulo the derived subgroup. *Arch. Math.* **56** (1991), 218–227.
- [2] G. Endimioni. Applications rationnelles d'un groupe nilpotent. *C. R. Acad. Sci. Paris* **314** (1992), 431–434.
- [3] G. Endimioni. Pointwise inner automorphisms in a free nilpotent group. *Quart. J. Math.* **53** (2002), 397–402.
- [4] G. Endimioni. On the polynomial automorphisms of a group. *Acta Sci. Math. (Szeged)* **73** (2007), 61–69.
- [5] S. Franciosi and F. de Giovanni. On automorphisms fixing normal subgroups of nilpotent groups. *Boll. Un. Mat. Ital. B* **7** (1987), 1161–1170.
- [6] C.K. Gupta. IA-automorphisms of two-generator metabelian groups. *Arch. Math.* **37** (1981), 106–112.

- [7] Ju. V. Kuz'min. Inner endomorphisms of metabelian groups (Russian). *Sibirsk. Mat. Zh.* **16** (1975), 736–744; translation in *Siberian Math. J.* **16** (1976), 563–568.
- [8] A. Lubotzky. Normal automorphisms of free groups. *J. Algebra* **63** (1980), 494–498.
- [9] H. Neumann. *Varieties of groups* (Springer-Verlag, 1967).
- [10] D.J.S. Robinson. *A course in the theory of groups* (Springer-Verlag, 1996).
- [11] N.S. Romanovskiĭ. Normal automorphisms of free solvable pro- p -groups (Russian). *Algebra i Logika* **36** (1997), 441–453; translation in *Algebra and Logic* **36** (1997), 257–263.
- [12] D. Schweigert. Polynomautomorphismen auf endlichen Gruppen. *Arch. Math.* **29** (1977), 34–38.
- [13] D. Segal. *Polycyclic groups* (Cambridge University Press, 1983).

C.M.I-UNIVERSITÉ DE PROVENCE, 39, RUE F. JOLIOT-CURIE, F-13453 MARSEILLE CEDEX 13

E-mail address: endimion@gypsis.univ-mrs.fr